
Mental Health Current Awareness Update

July 2022

Welcome to the monthly **Mental Health Current Awareness Alert** (MHCAU), produced by **UKHSA Knowledge and Library Services**. The purpose of this update is to provide you with the latest, best evidence about mental health, to help you make informed decisions.

If you have any queries, please contact us at libraries@phe.gov.uk, and if you would like to request a literature search, please use the **Knowledge and Library Services Enquiries Portal**.

Please note that not all the articles and resources referred to in this alert are freely available. Some articles may require an Athens username and password. For further information on Athens accounts please visit [here](#). If you are not eligible to access our library services, then you may want to approach your **local health library service**.

We do not accept responsibility for the availability or reliability of the items of content included in this alert and their inclusion is not an endorsement of any views that may be expressed.

-
- **Mental health and COVID-19 - 3**
 - **Mental health - 52**
 - **Prevention and mental well-being - 4**
 - **Psychiatry - 4**
 - **Depression - 36**
 - **Anxiety - 6**
 - **Personality disorder - 18**
 - **Psychosis - 4**
 - **Schizophrenia - 6**
 - **Eating disorders - 1**
 - **Self-harm - 5**
 - **Suicide - 12**
 - **Attention Deficit Hyperactivity Disorder - 1**
 - **Obsessive Compulsive Disorder - 1**

Mental health and COVID-19

COVID-19 ARDS and Posttraumatic Stress Disorder in Family Members After ICU Discharge—Reply

19 Jul 2022 - The JAMA Network

In Reply Among family members of patients with COVID-19 ARDS included in our study, half were indeed spouses. Dr Lin and colleagues suggest that as these family members were quarantined, this might have been an additional source of PTSD. However, recent findings seem to challenge this hypothesis

COVID-19 ARDS and Posttraumatic Stress Disorder in Family Members After ICU Discharge

19 Jul 2022 - The JAMA Network

To the Editor We would like to make several points that were not highlighted in the study about the association of COVID-19–related acute respiratory distress syndrome (ARDS) with symptoms of posttraumatic stress disorder (PTSD) in family members after intensive care unit (ICU) discharge.

RT @LancetRH_Europe: Our July Issue is online! Highlights include :

✔ Editorial: Lessons from #COVID19 are shaping the response to the monkeypox outbreak ✔ Stress-related disorders & mortality ✔ Call for action to adopt integrated care models to manage Post-Covid-19 Condition <https://t.co/ZM7u3LsE82> <https://t.co/XvTF5ZKxTn>

01 Jul 2022 - Twitter

RT @LancetRH_Europe: Our July Issue is online! Highlights include : ✔ Editorial: Lessons from #COVID19 are shaping the response to the monkeypox outbreak ✔ Stress-related disorders & mortality ✔ Call for action to adopt integrated care models to manage Post-Covid-19 Condition <https://www.thelancet.com/journals/lanep/home>

[\[back to top\]](#)

Mental health

A workplace organisational intervention to improve hospital nurses and physicians mental health: study protocol for the Magnet4Europe wait list cluster randomised controlled trial

28 Jul 2022 - British Medical Journal

Introduction The increasing burden of mental distress reported by healthcare professionals is a matter of serious concern and there is a growing recognition of the role of the workplace in creating this problem. Magnet hospitals, a model shown to attract and retain staff in US research, creates positive work environments that aim to support the well-being of healthcare professionals

Using prosocial behavior to safeguard mental health and foster emotional well-being during the COVID-19 pandemic: A registered report of a randomized trial

28 Jul 2022 - PLOS ONE

= 340). Findings Those in the prosocial condition did not differ on any outcome from those in the self-focused or neutral acts conditions during the intervention or at follow-up, nor did prosocial effects differ for those who had been negatively affected socially or economically by the pandemic (all p's > 0.05)

Policies on mental health in the workplace during the COVID-19 pandemic: A scoping review

28 Jul 2022 - PLOS ONE

by David Villarreal-Zegarra, C. Mahony Reátegui-Rivera, Iselle Sabastizagal-Vela, Miguel Angel Burgos-Flores, Nieves Alejandra Cama-Ttito, Jaime Rosales-Rimache Background The COVID-19 pandemic has had a profound impact on both mental health and working conditions. Workplaces are conducive spaces for implementing strategies and interventions to promote mental health

Impact on patient-provider relationship and documentation practices when mental health patients access their electronic health records

online: a qualitative study among health professionals in an outpatient setting

28 Jul 2022 - Biomed Central

Patient accessible electronic health records (PAEHR) hold the potential to increase patient empowerment, especially for patients with complex, long-term or chronic conditions. However, evidence of its benefits...

Gendered impact of COVID-19 containment measures on unpaid care work and mental health in Europe: a scoping review protocol

27 Jul 2022 - British Medical Journal

Introduction Women are more likely than men to provide unpaid care work. Previous research has shown that lack of support for various forms of unpaid care work and work-family conflicts have negative impacts on caregivers' mental health, especially among female caregivers

Role of the regulator in enabling a just culture: a qualitative study in mental health and hospital care

27 Jul 2022 - British Medical Journal

Objectives A just culture is considered a promising way to improve patient safety and working conditions in the healthcare sector, and as such is also of relevance to healthcare regulators who are tasked with monitoring and overseeing quality and safety of care

Assessing an Internet-Delivered, Emotion-Focused Intervention Compared With a Healthy Lifestyle Active Control Intervention in Improving Mental Health in Cancer Survivors: Protocol for a Randomized Controlled Trial

27 Jul 2022 - National Institutes of Health (NIH)

CONCLUSIONS: This is one of the first trials to develop and investigate the efficacy of a web-based intervention for cancer survivors that specifically targets emotion regulation.

Russian adolescent mental health in 2002, 2015 and during the COVID-19 pandemic in 2021

27 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Design of and outcomes in a student-run free mental health clinic serving the uninsured in East Harlem

26 Jul 2022 - Biomed Central

Safety-net clinics are an important source of low-cost or free mental healthcare to those with limited financial resources. Such clinics are often staffed by trainees in early stages of their career. Only limi...

Effectiveness and active ingredients of social prescribing interventions targeting mental health: a systematic review

25 Jul 2022 - British Medical Journal

Objective This study aims to establish the effectiveness and active ingredients of UK-based social prescribing interventions targeting mental health and well-being outcomes. Design Systematic review adhering to Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines and a published protocol. Data sources Nine databases were systematically searched up to March 2022

Parent and child mental health during COVID-19 in Australia: The role of pet attachment

25 Jul 2022 - PLOS ONE

by Shannon K. Bennetts, Sharinne B. Crawford, Tiffani J. Howell, Fiona Burgemeister, Catherine Chamberlain, Kylie Burke, Jan M. Nicholson Restrictions, social isolation, and uncertainty related to the global COVID-19 pandemic have disrupted the ways that parents and children maintain family routines, health, and wellbeing

Mental health indicators and their lifestyle associations in German students: a gender-specific multivariable analysis

25 Jul 2022 - Biomed Central

Mental health and wellbeing at work

25 Jul 2022 - NHS Networks

This session is designed to give insight and understanding to managers within the healthcare sector around how to support the people they manage, in identifying and addressing mental health concerns...

Serious Mental Illness in Healthcare and Academia: A Lived Experience

25 Jul 2022 - John Wiley & Sons

Journal of Psychiatric and Mental Health Nursing, Accepted Article.

Conceptualising public mental health: development of a conceptual framework for public mental health

24 Jul 2022 - Biomed Central

Numerous determinants have been linked to public mental health; however, they have not been brought together in a comprehensive conceptual framework. The goal of this work was to bring together academic research...

Incidence and associations of hospital delirium diagnoses in 85979 people with severe mental illness: a data linkage study

23 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

Prospective cohort study of the evaluation of patient benefit from the redevelopment of a complete national forensic mental health service: the Dundrum Forensic Redevelopment Evaluation Study (D-FOREST) protocol

22 Jul 2022 - British Medical Journal

Introduction Secure forensic mental health services are low volume, high cost services. They offer care and treatment to mentally disordered offenders who pose a high risk of serious violence to others. It is therefore incumbent on these services to systematically evaluate the outcomes of the care and treatment they deliver to ensure patient benefit in multiple domains

A mirror-image analysis of psychiatric hospitalisations among people with severe mental illness using Independent Supported Housing

22 Jul 2022 - Biomed Central

Evidence on the effectiveness of Independent Supported Housing (ISH) for non-homeless people with severe mental illness primarily comes from observational cohort studies, which have high risk of bias due to co...

[Articles] The temporal relationship between severe mental illness diagnosis and chronic physical comorbidity: a UK primary care cohort study of disease burden over 10 years

21 Jul 2022 - The Lancet

Elevated odds of multiple conditions at the point of severe mental illness diagnosis suggest that early intervention on physical health parameters is necessary to reduce morbidity and premature mortality. Some physical conditions might be under-recorded in patients with schizophrenia relative to patients with other severe mental illness subtypes.

Development and validation of a nomogram for predicting the risk of mental health problems of factory workers and miners

21 Jul 2022 - British Medical Journal

Objective A nomogram for predicting the risk of mental health problems was established in a population of factory workers and miners, in order to quickly calculate the probability of a worker suffering from mental health problems. Methods A cross-sectional survey of 7500 factory workers and miners in Urumqi was conducted by means of an electronic questionnaire using cluster sampling method

Occupational mental health

21 Jul 2022 - NHS Networks

Being in a productive and supportive work environment is linked to better mental health. However, those experiencing mental health problems are often either excluded from the workplace or not supported...

Does medical disparity exist while treating severe mental illness patients with acute appendicitis in emergency departments? A real-world database study

21 Jul 2022 - Biomed Central

Patients with severe mental illness (SMI) have a shorter life expectancy and have been considered by the World Health Organization (WHO) as a vulnerable group. As the causes for this mortality gap are complex,...

Spanish Mental Health Nurses' experiences of Mechanical Restraint: A qualitative descriptive study

20 Jul 2022 - John Wiley & Sons

Journal of Psychiatric and Mental Health Nursing, Accepted Article.

'More than just numbers on a page?' A qualitative exploration of the use of data collection and feedback in youth mental health services

20 Jul 2022 - PLOS ONE

by Craig Hamilton, Kate Filia, Sian Lloyd, Sophie Prober, Eilidh Duncan Objectives This study aimed to explore current data collection and feedback practice, in the form of monitoring and evaluation, among youth mental health (YMH) services and healthcare commissioners; and to identify barriers and enablers to this practice

Improving children and young people's mental health services

20 Jul 2022 - NHS Networks

This briefing presents analysis from The Health Foundation's Networked Data Lab about children and young people's mental health.

Online Racism and Mental Health Among Black American Adolescents in 2020

19 Jul 2022 - Elsevier ScienceDirect

Publication date: Available online 19 July 2022 Source: Journal of the American Academy of Child & Adolescent Psychiatry Author(s): Juan Del Toro, Ming-Te Wang

The impact of cognitive impairment of individuals with Parkinson's disease on their caregivers' mental health: A systematic review protocol

19 Jul 2022 - PLOS ONE

by Paulina Beata Golińska, Łucja Bieleninik, Michał Harciarek, Mariola Bidzan Introduction Parkinson's disease is a motor disease, the second most common neurodegenerative disorder with cardinal symptoms including bradykinesia, rigidity, and rest tremor accompanied by cognitive difficulties

The lived experience of severe mental illness and long-term conditions: a qualitative exploration of service user, carer, and healthcare professional perspectives on self-managing co-existing mental and physical conditions

19 Jul 2022 - Biomed Central

People with severe mental illness (SMI), such as schizophrenia, have higher rates of physical long-term conditions (LTCs), poorer health outcomes, and shorter life expectancy compared with the general populati...

Developing good practice indicators to assist mental health practitioners to converse with young people about their online activities and impact on mental health: a two-panel mixed-methods Delphi study

19 Jul 2022 - Biomed Central

Online activity has been linked to poor mental health in children and young people, particularly those with existing vulnerability who may inadvertently or otherwise access harmful content. It is suggested hea...

Factors associated with poor mental health outcomes in nurses in COVID-19-designated hospitals in the postepidemic period in Guangdong Province: a cross-sectional study

18 Jul 2022 - British Medical Journal

Objective The early days of the COVID-19 pandemic placed enormous pressure and subsequent negative psychological problems on nurses, but at this stage of the year-long COVID-19 outbreak, the level of stress and negative emotions that nurses experience is unclear. Our study attempted to assess the factors influencing mental health status in nurses during the postepidemic period of COVID-19

Debate: Conduct disorder and the segregation of child mental health

16 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

[Comment] HIV, substance use, and mental health care in prisons

14 Jul 2022 - The Lancet

Incarceration, psychiatric and substance-use disorders, and infectious diseases form a syndemic.¹ That is, criminal justice systems gather people with pre-existing, serious health problems and exacerbate them

Effectiveness of universal school-based mindfulness training compared with normal school provision on teacher mental health and school climate: results of the MYRIAD cluster randomised controlled trial

12 Jul 2022 - British Medical Journal

Background Education is broader than academic teaching. It includes teaching students social–emotional skills both directly and indirectly through a positive school climate. Objective To evaluate if a universal school-based mindfulness training (SBMT) enhances teacher mental health and school climate

The impact of mindfulness training in early adolescence on affective executive control, and on later mental health during the COVID-19 pandemic: a randomised controlled trial

12 Jul 2022 - British Medical Journal

Background Previous research suggests that mindfulness training (MT) appears effective at improving mental health in young people. MT is proposed to work through improving executive control in affectively laden contexts. However, it is unclear whether MT improves such control in young people

Do mindfulness-based programmes improve the cognitive skills, behaviour and mental health of children and adolescents? An updated meta-analysis of randomised controlled trials

12 Jul 2022 - British Medical Journal

Question Mindfulness-based programmes (MBPs) are an increasingly popular approach to improving mental health in young people. Our previous meta-analysis suggested that MBPs show promising effectiveness, but highlighted a lack of high-quality, adequately powered randomised controlled trials (RCTs)

Effectiveness and cost-effectiveness of universal school-based mindfulness training compared with normal school provision in reducing risk of mental health problems and promoting well-being in adolescence: the MYRIAD cluster randomised controlled trial

12 Jul 2022 - British Medical Journal

Background Systematic reviews suggest school-based mindfulness training (SBMT) shows promise in promoting student mental health. Objective The My Resilience in Adolescence (MYRIAD) Trial evaluated the effectiveness and cost-effectiveness of SBMT compared with teaching-as-usual (TAU). Methods MYRIAD was a parallel group, cluster-randomised controlled trial

How can we optimise learning from trials in child and adolescent mental health?

12 Jul 2022 - British Medical Journal

Improving child and adolescent mental health requires the careful development and rigorous testing of interventions and delivery methods. This includes universal school-based mindfulness training, evaluated in the My Resilience in Adolescence (MYRIAD) trial reported in this special edition

Suicide following hospital admission for mental health conditions, physical illness, injury and intentional self-harm in Victoria, Australia

11 Jul 2022 - PLOS ONE

by Dasamal Tharanga Fernando, Angela Clapperton, Janneke Berecki-Gisolf Objective The majority of suicide decedents have had contact with health services close to their death. Some of these contacts include admissions to hospitals for physical and mental health conditions, injury and intentional self-harm

Parent and/or Grandparent Attendance at Residential School and Dimensions of Cultural Identity and Engagement: Associations with Mental Health and Substance Use among First Nations Adults Living off Reserve

11 Jul 2022 - Sage Journals

Society and Mental Health, Ahead of Print. Limited studies have assessed how parent and/or grandparent attendance at residential schools is associated with mental health and substance use among First Nations peoples living off reserve, while also considering how cultural dimensions relate to these outcomes

Preliminary Real-World Evidence Supporting the Efficacy of a Remote Neurofeedback System in Improving Mental Health: Retrospective Single-Group Pretest-Posttest Study

08 Jul 2022 - National Institutes of Health (NIH)

CONCLUSIONS: Preliminary findings suggest the efficacy of app-based remote neurofeedback in improving mental health, given the reduced symptom severity from pre- to postassessment for general psychological health, ADHD, anxiety, and depression, as well as adjusted resting EEG neural markers for individuals with symptoms of ADHD

Have the COVID-19 pandemic and lockdown affected childrens mental health in the long term? A repeated cross-sectional study

07 Jul 2022 - British Medical Journal

Objective The study aimed to evaluate the impact of the COVID-19 pandemic on levels of anxiety and depressive symptoms in children and adolescents. Design Cross-sectional surveys were carried out on the mental health of children; one survey was conducted before the COVID-19 pandemic and one into the pandemic, 15 months after the school closures and implementation of lockdown and social

Flooding and public mental health

07 Jul 2022 - NHS Networks

This guidance on flooding and health aims to provide information for public health authorities, emergency services and health professionals on the measures needed to prevent major effects on health arising...

Celebrating a lifetime award for a once controversial mental health nurse

07 Jul 2022 - RCNi

A once controversial figure in mental health nursing has received a major lifetime achievement award. Emeritus professor Kevin Gournay was given the Eileen Skellern Lifetime Achievement award for his contribution to the profession.

Nursing students' experiences of seeking mental health support: a literature review

07 Jul 2022 - RCNi

Nursing students are at risk of suboptimal mental health due to factors that are common to all students and factors that are specific to nursing courses. Supporting nursing students' mental health is important not only for each student's well-being but also for the nursing workforce as a whole. A systematic

Centering Culture in Mental Health: Differences in Diagnosis, Treatment, and Access to Care Among Older People of Color

07 Jul 2022 - Elsevier

In 2003, the Institute of Medicine (IOM) published Unequal Treatment, which elevated racial and ethnic healthcare disparities to the forefront of clinical and policy attention.¹ However,

since its publication, little progress has been made to lessen the disparities in mental healthcare, especially among older adults of color

Effectiveness of emotion focused skills training for parents: study protocol for a randomized controlled trial in specialist mental health care

07 Jul 2022 - Biomed Central

Emotion-Focused Skills Training (EFST) is a newly developed manualized skill training program for parents to strengthen emotional bonds between parents and children and improve mental health outcomes in childr...

"I lost so much more than my partner" – Bereaved partners' grief experiences following suicide or physician-assisted dying in case of a mental disorder

07 Jul 2022 - Biomed Central

There is a lack of existing research on grief following the intentional death of people suffering from a mental disorder. Our study aims to provide insight into grief experiences and social reactions of bereav...

Neighbourhood and family correlates of immigrant children's mental health: a population-based cross-sectional study in Canada

05 Jul 2022 - Biomed Central

Immigrant children exhibit significant variation in their mental health outcomes despite disproportionate exposure to socio-economic adversity compared to their non-immigrant peers. Identifying aspects of neig...

Social networks and internet emotional relationships on mental health and quality of life in students: structural equation modelling

05 Jul 2022 - Biomed Central

Social networks and relationships create a sense of belonging and social identity; hence, can affect mental health and the quality of life, especially in young people. The present study was conducted to determ...

A cluster randomized trial for improving mental health and well-being of persons affected by leprosy or buruli ulcer in Nigeria: A study protocol

01 Jul 2022 - National Institutes of Health (NIH)

This protocol describes a study in which we would assess the effect of using community lay counselors, self-help groups (SHGs), and trained frontline health workers to reduce mental disorders and improve quality of life (QOL) of persons affected by leprosy or Buruli ulcer (BU). A cluster randomized controlled study design will be employed. The study will involve persons affected by leprosy or BU

Mental health services in response to the COVID-19 pandemic in high-income countries: protocol for a rapid review

30 Jun 2022 - British Medical Journal

Introduction The COVID-19 pandemic has caused disruptions to mental health services, forcing the rapid implementation of alternative ways of delivering services alongside a greater immediate, and continuously growing, demand across those services

Mental health and working constraints of first-year PhD students in health and science in a French university: a cross-sectional study in the context of occupational health monitoring

30 Jun 2022 - British Medical Journal

Objectives To assess the mental health and working constraints perception of first-year PhD in France. Design It is a cross-sectional study. Setting The study was conducted by the team of the Lyon 1 University's Occupational Health and Medical Service for Staff, from October 2019 to February 2020

[\[back to top\]](#)

Prevention and mental well-being

Prevalence of Health-Risk Behaviors and Mental Well-Being of ASEAN University Students in COVID-19 Pandemic

27 Jul 2022 - National Institutes of Health (NIH)

The prevalence of epidemiological health-risk behaviors and mental well-being in the COVID-19 pandemic, stratified by sociodemographic factors in Association of South East Asian Nations (ASEAN) university students, were examined in the research. Data were collected in March-June 2021 via an online survey from 15,366 university students from 17 universities in seven ASEAN countries

Neonatal nurse practitioner job satisfaction, workforce environment, and mental well-being

06 Jul 2022 - National Institutes of Health (NIH)

CONCLUSIONS: Neonatal nurse practitioners' satisfaction is multifactorial. Those who reported taking time off for self-prescribed mental health indicated less job satisfaction, more work hours, poor work/life balance, and a less-than-optimum work environment.

TWIN-10: protocol for a 10-year longitudinal twin study of the neuroscience of mental well-being and resilience

01 Jul 2022 - British Medical Journal

Introduction Mental well-being is a core component of mental health, and resilience is a key process of positive adaptive recovery following adversity. However, we lack an understanding of the neural mechanisms that contribute to individual variation in the trajectories of well-being and resilience relative to risk. Genetic and/or environmental factors may also modulate these mechanisms

Student Experience and Its Relationship with Campus Mental Well-Being

01 Jul 2022 - National Institutes of Health (NIH)

This research offers a generalizable Campus Mental Well-being Sense of Coherence Framework for improving student experience by classifying SES variables according to Antonovsky's salutogenic health logic (GRRs and SRRs) and by mapping these variables to the Information Infrastructure to Experience Framework (IEF).

[\[back to top\]](#)

Psychiatry

The Enactment Concept in Child and Adolescent Psychiatry Literature

15 Jul 2022 - Elsevier ScienceDirect

Publication date: Available online 15 July 2022Source: Journal of the American Academy of Child & Adolescent PsychiatryAuthor(s): Timothy Rice, Rachel Z. Ritvo

[Personal View] More phenomenology in psychiatry? Applied ontology as a method towards integration

08 Jul 2022 - The Lancet

There have been renewed calls to use phenomenology in psychiatry to improve knowledge about causation, diagnostics, and treatment of mental health conditions. A phenomenological approach aims to elucidate the subjective experiences of mental health, which its advocates claim have been largely neglected by current diagnostic frameworks in psychiatry (eg, DSM-5)

[Corrections] Correction to Lancet Psychiatry 2022; 9: e32

06 Jul 2022 - The Lancet

Cacciatore J, Frances A. DSM-5-TR turns normal grief into a mental disorder. Lancet Psychiatry 2022; 9: e32—In this Correspondence, the spelling of Allen Frances's name was incorrect. This correction has been made to the online version as of July 6, 2022.

JAMA Psychiatry

01 Jul 2022 - The JAMA Network

Mission Statement: JAMA Psychiatry strives to publish original, state-of-the-art studies and commentaries of general interest to clinicians, scholars, and research scientists in psychiatry, mental health, behavioral science, and allied fields

[\[back to top\]](#)

Depression

Predictors of longer-term depression trajectories during the COVID-19 pandemic: a longitudinal study in four UK cohorts

28 Jul 2022 - British Medical Journal

BackgroundThe COVID-19 pandemic has caused an increase in mental ill health compared with prepandemic levels. Longer-term trajectories of depression in adults during the pandemic remain unclear.**Objective**We used latent growth curve modelling to examine individual trajectories of depression symptoms, and their predictors, beyond the early stage of the pandemic

Treatment options for depression

28 Jul 2022 - NHS Networks

NICE's updated guideline on the treatment and management of depression in adults advises that patients should make an informed choice on which treatment option is right for them.

Information needs and sources of information among people with depression and anxiety: a scoping review

27 Jul 2022 - Biomed Central

Previous studies have identified substantial unmet information needs in people with depression and anxiety. Sufficient information about the disorder, treatment, available services, and strategies for self-man...

Sustained remission from perinatal depression after bright light therapy: a pilot randomised, placebo-controlled trial

25 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

School Performance and Later Diagnoses of Non-Affective Psychoses, Bipolar Disorder and Depression

25 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

Depression and anxiety and their associated factors among caregivers of children and adolescents with epilepsy in three selected hospitals in Amhara region, Ethiopia: A cross-sectional study

25 Jul 2022 - PLOS ONE

by Mekonnen Tsehay, Mogesie Necho, Asmare Belete, Mengesha Srahbzu **Background** The prevalence rates of depression and anxiety are unforeseen among primary caregivers of patients with epilepsy. Little attention is being given to the problem in Ethiopia

Effects of insomnia and levels of depression and anxiety symptoms on quality of life in people with epilepsy

25 Jul 2022 - Biomed Central

The association between insomnia and quality of life (QOL) in epilepsy is poorly understood and may involve interactive variables. We aimed to investigate whether and how insomnia, levels of depression and anx...

A Southeast Asian expert consensus on the management of major depressive disorder with suicidal behavior in adults under 65 years of age

21 Jul 2022 - Biomed Central

The high prevalence of suicidal behavior among individuals with major depressive disorder (MDD) in Southeast Asia (SEA) underscores the need for optimized management to address depressive symptoms, reduce suic...

Association between life events and later depression in the population-based Heinz Nixdorf Recall study—The role of sex and optimism

20 Jul 2022 - PLOS ONE

by Janine Gronewold, Ela-Emsal Duman, Miriam Engel, Miriam Engels, Johannes Siegrist, Raimund Erbel, K-H. Jöckel, Dirk M. Hermann **Background** The association between life

event stress and depressive symptoms has not been analyzed in the general population before. Methods In the population-based Heinz Nixdorf Recall study, we assessed the association of 1

Effectiveness of interventions to improve medication adherence in adults with depressive disorders: a meta-analysis

20 Jul 2022 - Biomed Central

Non-adherence to medication is a major obstacle in the treatment of depressive disorders. We systematically reviewed the literature to evaluate the effectiveness of interventions aimed at improving adherence t...

Feasibility and acceptability of a novel telepsychiatry-delivered precision prescribing intervention for anxiety and depression

19 Jul 2022 - Biomed Central

Major Depressive Disorder and Generalized Anxiety Disorder are pervasive and debilitating conditions, though treatment is often inaccessible and based on trial-and-error prescribing methods. The present observ...

Appearance satisfaction mediates the relationship between recreational screen time and depressive symptoms in adolescents

18 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Web-based treatment for depression in pregnancy: a feasibility study of Mum2BMoodBooster

16 Jul 2022 - Biomed Central

Depression in pregnancy is prevalent, under-treated, and has serious impacts on the wellbeing of women and on child development. Internet programs can reach women who may not access traditional treatments due ...

Clinical practice guidelines with recommendations for peripartum depression: A European systematic review

15 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

Psychomotor semiology in depression: a standardized clinical psychomotor approach

15 Jul 2022 - Biomed Central

Although psychomotor symptoms are associated with the clinical symptomatology of depression, they are rarely assessed and standardized clinical evaluation tools are lacking. Psychomotor retardation is sometime...

Health care professionals' experiences of screening immigrant mothers for postpartum depression—a qualitative systematic review

14 Jul 2022 - PLOS ONE

by Malin Skoog, Inger Kristensson Hallström, Andreas Vilhelmsson Introduction Postpartum depression is considered a major public health problem, which immigrant mothers are at particular risk of being affected by, but it can also have long-lasting traumatic effects on the child's health and development

University students' help seeking intention for depression from health professionals; a cross sectional study

13 Jul 2022 - PLOS ONE

by Alemu Lemma, Woredaw Minichil, Endalamaw Salelew, Jinenus Tadesa, Habtamu Kerebih, Kabtamu Nigussie, Demeke Demilew, Shegaye Shumet Objectives To assess University students' intention to seek help for depression from health professional and associated factors among University of Gondar students, Northwest, Ethiopia

Universal prevention of depression at schools: dead end or challenging crossroad?

12 Jul 2022 - British Medical Journal

Universal school programmes aimed at the prevention of depression and other common mental health problems in adolescents are attractive because they are less stigmatising than

targeted interventions, have a high uptake and may shift the 'normal distribution' of mental health problems in the positive direction

Modular-based psychotherapy (MoBa) versus cognitive-behavioural therapy (CBT) for patients with depression, comorbidities and a history of childhood maltreatment: study protocol for a randomised controlled feasibility trial

12 Jul 2022 - British Medical Journal

Introduction In depression treatment, most patients do not reach response or remission with current psychotherapeutic approaches. Major reasons for individual non-response are interindividual heterogeneity of etiological mechanisms and pathological forms, and a high rate of comorbid disorders

Moderate-to-vigorous group aerobic exercise versus group leisure activities for mild-to-moderate depression in adolescents: study protocol for a multicentre randomised controlled trial

12 Jul 2022 - British Medical Journal

Introduction Depression is common, increasing among adolescents and carries risk of disability, lower educational achievements, cardiovascular disease, substance abuse, self-harm and suicide. The effects of evidence-based treatments with medication or psychotherapy are modest

Habenula activation patterns in a preclinical model of neuropathic pain accompanied by depressive-like behaviour

12 Jul 2022 - PLOS ONE

by Geiza Fernanda Antunes, Ana Carolina Pinheiro Campos, Danielle Varin de Assis, Flavia Venetucci Gouveia, Midiã Dias de Jesus Seno, Rosana Lima Pagano, Raquel Chacon Ruiz Martinez Pain and depression are complex disorders that frequently co-occur, resulting in diminished quality of life

Examining academic self-concept as a mediator of the relationship between anxiety and depression: A longitudinal study

11 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Rapid-Response Treatments for Depression and Requests for Physician-Assisted Death: An Ethical Analysis

10 Jul 2022 - Elsevier

Depression is common at the end of life, and there is longstanding concern that it may affect terminally ill patients' decisions to request physician-assisted death (PAD). However, it is difficult for clinicians to determine the role of depression in a patient's PAD request

Prevalence of and risk factors for depression, anxiety, and stress in non-hospitalized asymptomatic and mild COVID-19 patients in East Java province, Indonesia

07 Jul 2022 - PLOS ONE

by Michael Austin Pradipta Lusida, Sovia Salamah, Michael Jonatan, Illona Okvita Wiyogo, Claudia Herda Asyari, Nurarifah Destianizar Ali, Jose Asmara, Ria Indah Wahyuningtyas, Erwin Astha Triyono, Ni Kadek Ratnadewi, Aryan Irzaldy, Firas Farisi Alkaff Background Despite abundant data on mental health during the COVID-19 pandemic, 3 important knowledge gaps continue to exist, i.e

Associations between WASH-related violence and depressive symptoms in adolescent girls and young women in South Africa (HPTN 068): a cross-sectional analysis

05 Jul 2022 - British Medical Journal

Objective There is a lack of research on experiences of WASH-related violence. This study aims to quantify the association between experience or worry of violence when using the toilet or collecting water and depressive symptoms among a cohort of young women in South Africa. Methods Data are from visit 3 of the HPTN 068 cohort of adolescent girls in rural Mpumalanga Province, South Africa

Exploring bridge symptoms in HIV-positive people with comorbid depressive and anxiety disorders

05 Jul 2022 - Biomed Central

The prevalence of comorbid depressive and anxiety disorders in people living with HIV (PLWH) is high. However, it is unclear which symptom is the bridge symptom between depression and anxiety in PLWH. This stu...

Prevalence and correlates of REM sleep behaviour disorder in patients with major depressive disorder: a two-phase study

04 Jul 2022 - British Medical Journal

ObjectiveTo investigate the prevalence and clinical correlates of video polysomnography (vPSG)-confirmed rapid eye movement sleep behaviour disorder (RBD) in patients with major depressive disorder (MDD). **Methods**This is a clinic-based two-phase epidemiological study. In phase 1, patients with MDD were screened by a validated questionnaire, RBD Questionnaire-Hong Kong (RBDQ-HK)

Five-year longitudinal evidence supports the safety and efficacy of electroconvulsive therapy for older adults with major depressive disorder

04 Jul 2022 - Elsevier

Major depressive disorder (MDD) in older adults is a severe neuropsychiatric illness that results in significant disability, morbidity, and mortality¹. While antidepressant treatment strategies have evolved considerably in recent years, many may be ineffective and unable to alleviate depressive symptoms²

Prevalence and associated factors depressive symptoms among healthcare professionals at Dessie comprehensive specialized hospital, Ethiopia

04 Jul 2022 - Biomed Central

Depression among healthcare professionals results in adverse effects which might include decreased impairment of work performance, an increased turnover rate, and vulnerability to clinical error. Despite that,...

Symptoms of depression, anxiety and stress in patients with chronic otitis media

01 Jul 2022 - PLOS ONE

by Ana D. Jotic, Ana M. Opankovic, Zorana Z. Radin, Ljiljana Cvorovic, Katarina R. Savic Vujovic, Sanja B. Krejovic-Trivic, Bojana M. Bukurov, Biljana R. Milicic, Jasmina D. Stojanovic **Purpose** Persistent symptoms of chronic otitis media cause limitations in daily routine and social interactions, influencing significantly patients' quality of life and mental health

Evaluation of Depression Severity With Ketamine and Esketamine Treatment in a Clinical Setting

01 Jul 2022 - The JAMA Network

This comparative analysis examines the trajectory of depression severity among patients treated with intravenous ketamine or intranasal esketamine in a clinical setting.

Comorbidity and Coaggregation of Major Depressive and Bipolar Disorders and Cannabis Use Disorder

01 Jul 2022 - The JAMA Network

This cross-sectional study evaluates familial associations between cannabis use disorder and bipolar disorder II.

In the here and now: Future thinking and social problem-solving in depression

30 Jun 2022 - PLOS ONE

by Saima Noreen, Barbara Dritschel This research investigates whether thinking about the consequences of a problem being resolved can improve social problem-solving in clinical depression. We also explore whether impaired social problem solving is related to inhibitory control

Prevalence of probable depression and factors associated with mean Hopkins Symptom Checklist (HSCCL) depression score among young women at high risk aged 15–24 years in Kampala, Uganda

30 Jun 2022 - PLOS ONE

by Onesmus Kamacooko, Daniel Bagiire, Francis Xavier Kasujja, Miriam Mirembe, Janet Seeley, Rachel King Background In populations at high risk of HIV infection, rates of depression can be elevated with far-reaching effects on overall well-being. There is limited research on depression among young women engaged in high-risk sexual behaviour in low and middle-income settings

Study protocol for the Alzheimer and music therapy study: An RCT to compare the efficacy of music therapy and physical activity on brain plasticity, depressive symptoms, and cognitive decline, in a population with and at risk for Alzheimer's disease

30 Jun 2022 - PLOS ONE

by Birthe K. Flo, Anna Maria Matziorinis, Stavros Skouras, Tobba Therkildsen Sudmann, Christian Gold, Stefan Koelsch Background There is anecdotal evidence for beneficial effects of music therapy in patients with Alzheimer's Disease (AD). However, there is a lack of rigorous research investigating this issue

Depression, anxiety, and academic performance in COVID-19: a cross-sectional study

30 Jun 2022 - Biomed Central

Depression and anxiety are common after months of social isolation, and they can have a negative impact on anyone's quality of life if they are not treated promptly and appropriately. The aim of this study was...

[\[back to top\]](#)

Anxiety

Effects of group mindfulness-based cognitive therapy and group cognitive behavioural therapy on symptomatic generalized anxiety disorder: a randomized controlled noninferiority trial

19 Jul 2022 - Biomed Central

Mindfulness-based cognitive therapy (MBCT) is a promising alternative treatment for generalized anxiety disorder (GAD). The objective of this study was to examine whether the efficacy of group MBCT adapted for...

The association between psychological burden related to COVID-19 and addictive social media use: Testing the mediational role of anxious affect

12 Jul 2022 - PLOS ONE

by Zahir Vally, Mai Helmy Introduction The COVID-19 pandemic has had a substantial impact on the normalcy of life. Similarly, social media use (SMU) has increased exponentially. This study examined the association between individuals' perception of the psychological burden related to the pandemic and addictive SMU

Research shows around 2 in 5 parents of hospitalised babies have anxiety or post-traumatic stress in the first month after birth. These levels are much higher than those generally seen in new parents:

[@NPEU_Oxford #PRUMNHC @MineshKhashu @Blisscharity](https://t.co/jscKYBDECn) <https://t.co/LfQ3hLucJa>

12 Jul 2022 - Twitter

Research shows around 2 in 5 parents of hospitalised babies have anxiety or post-traumatic stress in the first month after birth. These levels are much higher than those generally seen in new parents: <https://evidence.nihr.ac.uk/alert/neonatal-care-parents-may-need-long-term-psychological-support/> [@NPEU_Oxford #PRUMNHC @MineshKhashu @Blisscharity](#)

Factors influencing self-management among patients diagnosed with anxiety disorders in China: A cross-sectional study

11 Jul 2022 - John Wiley & Sons

Journal of Psychiatric and Mental Health Nursing, Accepted Article.

Factor structure, measurement invariance, and concurrent validity of the Patient Health Questionnaire-9 and the Generalized Anxiety Disorder scale-7 in a Norwegian psychiatric outpatient sample

11 Jul 2022 - Biomed Central

The aim of this study was to test factor structure, measurement invariance, and concurrent validity of the nine item Patient Health Questionnaire-9 (PHQ-9) and the seven item Generalized Anxiety Disorder scale...

Around 2 in 5 (40%) parents have anxiety or post-traumatic stress when their newborn baby needs intensive care. Research found that levels declined over time, but a year after the birth, 1 in 4 (25%) were still suffering: [@NPEU_Oxford @Blisscharity](https://t.co/jscKYBDECn) <https://t.co/1FtdFOdhGc>

30 Jun 2022 - Twitter

Around 2 in 5 (40%) parents have anxiety or post-traumatic stress when their newborn baby needs intensive care. Research found that levels declined over time, but a year after the birth, 1 in 4 (25%) were still suffering: <https://evidence.nihr.ac.uk/alert/neonatal-care-parents-may-need-long-term-psychological-support/> @NPEU_Oxford @Blisscharity

[back to top]

Personality disorder

Commentary: Against the widespread use of other types of PD diagnosis – a commentary on the May 2022 debate (should CAMH professionals be diagnosing personality disorder in adolescence?)

27 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Commentary: Bringing together lived experience, clinical and research expertise – a commentary on the May 2022 debate (should CAMH professionals be diagnosing personality disorder in adolescence?)

23 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Response: Personality disorder is here to stay across the lifespan – a response to Commentaries on the May 2022 Debate

17 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

Childhood symptoms of Attention-Deficit/Hyperactivity Disorder and Borderline Personality Disorder

14 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

The use of borderline personality disorder severity index-iv feedback in adjusting borderline personality disorder treatment: therapists and patients perspectives

14 Jul 2022 - Biomed Central

Previous research has emphasized the importance of therapists giving Routine Outcome Monitoring (ROM) feedback to their patients. It has been shown that several factors influence therapists' tendency to provid...

The trait model of the DSM–5 alternative model of personality disorder (AMPD): A structural review.

04 Jul 2022 - American Psychological Association

Publication of the Diagnostic and Statistical Manual for Mental Disorders, Fifth Edition (American Psychiatric Association, 2013), which included an alternative model of personality disorders (AMPD), was followed by an explosion of publications on the new model

An agenda for establishing the DSM–5 alternative model of personality disorder (AMPD) trait model in adolescence: Comment on Clark and Watson (2022).

04 Jul 2022 - American Psychological Association

In the years since the publication of the Diagnostic and Statistical Manual for Mental Disorders, Fifth Edition (DSM–5; American Psychiatric Association, 2013), considerable progress has been made in understanding the structure of the traits included in the Alternative Model of Personality Disorders (AMPD) in adulthood

Sociocultural context and the DSM–5 alternative model of personality disorder: Comment on Widiger and Hines (2022).

04 Jul 2022 - American Psychological Association

Widiger and Hines (2022) provide a brief overview of the development of the alternative model of personality disorder (AMPD) housed within Section 3 of the Fifth Edition of the Diagnostic and Statistical Manual of Mental Disorders (American Psychiatric Association, 2013)

“Effects of diagnostic disclosure and varying diagnostic terminology on social attitudes to personality disorder: An experimental vignette study”: Correction to O’Connor and Murphy (2021).

04 Jul 2022 - American Psychological Association

Reports an error in "Effects of diagnostic disclosure and varying diagnostic terminology on social attitudes to personality disorder: An experimental vignette study" by Cliodhna O’Connor and Lisa Murphy (Personality Disorders: Theory, Research, and Treatment, 2021[May], Vol 12[3], 241-248). In the article, the online publication date was listed incorrectly

Antisocial personality disorder and psychopathy: The AMPD in review.

04 Jul 2022 - American Psychological Association

The current review examines the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM–5) alternative model for personality disorders’ (AMPD) operationalization of antisocial personality disorder (ASPD), particularly as it relates to the construct of psychopathy

Is the alternative model for personality disorders able to capture avoidant personality disorder according to Section II of the DSM–5? A systematic review.

04 Jul 2022 - American Psychological Association

This review aims at examining the continuity between the categorical model for personality disorders (PDs) as defined by Section II of the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, and the alternative model for personality disorders (AMPD) with respect to Section II avoidant personality disorder (AvPD)

Ten-year retrospective on the DSM–5 alternative model of personality disorder: Seeing the forest for the trees.

04 Jul 2022 - American Psychological Association

This is an introduction to the special issue “The DSM-5 Alternative Model of Personality Disorders: A Ten-Year Retrospective.” The authors begin by summarizing the main findings and recommendations from each target review and its commentaries, as well as the shorter review articles, after which they offer additional remarks in the hopes of contributing to the setting of an agenda for the next

The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition alternative model of personality disorder.

04 Jul 2022 - American Psychological Association

The fifth edition of the American Psychiatric Association’s Diagnostic and Statistical Manual of Mental Disorders (DSM–5; American Psychiatric Association, 2013) includes within a section for emerging measures and models an alternative model of personality disorder

(AMPD). This article provides a brief overview of its development, noting, in particular, issues and controversies

If an entrepreneur founded the DSM–5 alternative model of personality disorder: When Axis II meets disruptive innovation: Comment on Widiger and Hines (2022).

04 Jul 2022 - American Psychological Association

Personality disorder (PD) researchers proposed a highly innovative “paradigm-shifting” revamp for the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (APA, 2013). Yet, 10 years later, Widiger and Hines (2022) summarize a developmental process plagued by disagreement and stagnation, with little evidence of the field having reaped the desired benefits of this diagnostic

Schizotypal personality disorder in the alternative model for personality disorders.

04 Jul 2022 - American Psychological Association

Schizotypal personality disorder (SPD) first appeared in the American Psychiatric Association diagnostic nosology in 1980, although its roots stretch back more than 100 years under labels such as borderline, ambulatory, and latent schizophrenia

The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, alternative model conceptualization of borderline personality disorder: A review of the evidence.

04 Jul 2022 - American Psychological Association

In the 10 years following the publishing of the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM–5), studies have accumulated testing the validity and utility of the alternative model for personality disorders (AMPD) in the context of borderline personality disorder (BPD)

The alternative model of personality disorder is inadequate for capturing obsessive-compulsive personality disorder.

04 Jul 2022 - American Psychological Association

The alternative model of personality disorder (AMPD) included in Section III of the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM–5) reconceptualized personality pathology in terms of pathological traits and impairments in functioning

Narcissism and the DSM–5 alternative model of personality disorder.

26 May 2022 - American Psychological Association

In the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Section III alternative model of personality disorder, narcissistic personality disorder (NPD) is assessed via required elevations on 2 of 4 Criterion A domains of personality dysfunction (i.e., self-direction, identity, empathy, intimacy) and 2 personality traits (Criterion B: attention seeking and grandiosity)

[\[back to top\]](#)

Psychosis

Editorial: Psychotic-like Experiences: Bolstering Protective Factors in Marginalized Youth

14 Jul 2022 - Elsevier ScienceDirect

Publication date: Available online 14 July 2022Source: Journal of the American Academy of Child & Adolescent PsychiatryAuthor(s): Bernalyn Ruiz-Yu, Derek M. Novacek, Carrie E. Bearden

Understanding the inverse relationship between age and psychotic symptoms: the role of borderline personality traits

13 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

Clinical, Brain, and Multilevel Clustering in Early Psychosis and Affective Stages

01 Jul 2022 - The JAMA Network

This cohort study uses machine learning techniques to cluster, compare, and combine subgroup solutions using clinical and brain structural imaging data from early psychosis and depression stages.

The Course of General Cognitive Ability in Individuals With Psychotic Disorders

01 Jul 2022 - The JAMA Network

This cohort study investigates the course of cognition from elementary school to old age among individuals with schizophrenia and other psychotic disorders to test neurodevelopmental and neurodegenerative theories of psychotic disorders.

[\[back to top\]](#)

Schizophrenia

Influencing factors of multiple adverse outcomes among schizophrenia patients using count regression models: a cross-sectional study

15 Jul 2022 - Biomed Central

Schizophrenia patients have increased risks of adverse outcomes, including violent crime, aggressiveness, and suicide. However, studies of different adverse outcomes in schizophrenia patients are limited and t...

Unemployment, homelessness, and other societal outcomes in patients with schizophrenia: a real-world retrospective cohort study of the United States Veterans Health Administration database

08 Jul 2022 - Biomed Central

The burden associated with schizophrenia is substantial. Impacts on the individual, healthcare system, and society may be particularly striking within the veteran population due to the presence of physical and...

Health resource utilization and cost before versus after initiation of second-generation long-acting injectable antipsychotics among adults with schizophrenia in Alberta, Canada: a retrospective, observational single-arm study

02 Jul 2022 - Biomed Central

Long-acting injectable (LAI) antipsychotics, along with community treatment orders (CTOs), are used to improve treatment effectiveness through adherence among individuals with schizophrenia. Understanding real...

Exonic deletions in IMMP2L in schizophrenia with enhanced glycation stress subtype

01 Jul 2022 - PLOS ONE

by Akane Yoshikawa, Itaru Kushima, Mitsuhiro Miyashita, Kazuhiro Suzuki, Kyoka Iino, Kazuya Toriumi, Yasue Horiuchi, Hideya Kawaji, Norio Ozaki, Masanari Itokawa, Makoto Arai
We previously identified a subtype of schizophrenia (SCZ) characterized by increased plasma pentosidine, a marker of glycation and oxidative stress (PEN-SCZ)

Associating Violence With Schizophrenia—Risks and Biases

01 Jul 2022 - The JAMA Network

To the Editor Whiting et al conducted a systematic review and meta-analysis investigating the association between schizophrenia spectrum disorder and violence, including subgroup analysis by sex. They found that schizophrenia spectrum disorder was associated with a significant increase in risk of perpetrating violence compared with control, especially among women (odds ratio [OR], 10.2)

Associating Violence With Schizophrenia—Risks and Biases—Reply

01 Jul 2022 - The JAMA Network

In Reply In our systematic review and meta-analysis examining the association between schizophrenia spectrum disorders (SSD) and violence, we included 24 studies of violent outcomes. Data from 15 countries consistently reported associations between SSD and violent outcomes despite using different diagnostic criteria and violence measurement

[\[back to top\]](#)

Eating disorders

[Online eating disorder learning available](#)

07 Jul 2022 - NHS Networks

To support GPs and primary care clinicians in understanding and supporting people of all ages with an eating disorder, Health Education England has worked in partnership with experts by experience,...

[\[back to top\]](#)

Self-harm

[\[Articles\] Prevalence of suicidal ideation and self-harm behaviours in children aged 12 years and younger: a systematic review and meta-analysis](#)

27 Jul 2022 - The Lancet

A high number of children in the general population can experience suicidal ideation and self-harm behaviours, thus underlining the need for more research on childhood suicide, including developmentally appropriate preventive strategies, such as youth-nominated support teams or dialectical behavioural therapy.

[Self-Harm in Eating Disorders \(SHINE\): a mixed-methods exploratory study](#)

27 Jul 2022 - British Medical Journal

Introduction Self-harm is highly prevalent among young people with eating disorders. However, why a young person may develop and continue to experience both an eating disorder and self-harm is unclear. This study will investigate the frequency, intensity, duration, function, context and processes of self-harm among people aged 16–25 diagnosed with an eating disorder

[Autism and self-harm: a population-based and discordant sibling study of young individuals](#)

22 Jul 2022 - John Wiley & Sons

Acta Psychiatrica Scandinavica, Accepted Article.

[Digital self-harm and suicidality among adolescents](#)

11 Jul 2022 - John Wiley & Sons

Child and Adolescent Mental Health, EarlyView.

[Association of Psychiatric Care Following Treatment for Deliberate Self-harm With Prospective Mortality](#)

01 Jul 2022 - The JAMA Network

This cohort study evaluates the association between risk of death and follow-up psychiatric care among patients treated for deliberate self-harm.

[\[back to top\]](#)

Suicide

[Suicide ideation and/or attempt with substance use and associated factors among the youth in northwest Ethiopia, community-based](#)

28 Jul 2022 - Biomed Central

Substance use is referring to the use of psychoactive substances like chat, cigarettes, alcohol, and others. The use of substances particularly (alcohol, chat, and cigarette) is a major mental health burden in...

[Patient expectations of and experiences with a suicide risk identification algorithm in clinical practice](#)

23 Jul 2022 - Biomed Central

Suicide risk prediction models derived from electronic health records (EHR) and insurance claims are a novel innovation in suicide prevention but patient perspectives on their use have been understudied.

Metabolomic biomarkers related to non-suicidal self-injury in patients with bipolar disorder

22 Jul 2022 - Biomed Central

Non-suicidal self-injury (NSSI) is an important symptom of bipolar disorder (BD) and other mental disorders and has attracted the attention of researchers lately. It is of great significance to study the chara...

Editorial: Antibullying Initiatives and Suicide Prevention

19 Jul 2022 - Elsevier ScienceDirect

Publication date: Available online 19 July 2022Source: Journal of the American Academy of Child & Adolescent PsychiatryAuthor(s): Alice Charach

Trapped in my inner prison—Cross-sectional examination of internal and external entrapment, hopelessness and suicidal ideation

19 Jul 2022 - PLOS ONE

by Inken Höller, Amelie Kremers, Dajana Schreiber, Thomas Forkmann Background Within the integrated motivational-volitional model of suicidal behavior, entrapment that consecutively leads to hopelessness is considered as a proximal risk factor for suicidal ideation

Psychoactive substances use before suicide: detailed analysis of all cases that occurred in the Brazilian Federal District in a 10-years period

14 Jul 2022 - Biomed Central

Psychoactive substances (PASs) are an important risk factor for suicide. This study investigated the sociodemographic characteristics, data related to the suicidal behavior, the methods employed, the circumsta...

Magnitude and associated factors of suicidal behavior among postpartum mothers attending public health centers of Addis Ababa, Ethiopia

13 Jul 2022 - Biomed Central

In low-income nations, suicidal conduct increases within the first year following delivery, ranging from 4 to 17.6%, with a three-fold cause of maternal death. Suicidal behavior was also reported to be 14% amo...

Commentary on Lange et al.: Acute alcohol use before suicide—is it contributing to an increase in suicide rates in the United States?

12 Jul 2022 - John Wiley & Sons

Addiction, EarlyView.

A state level analyses of suicide and the COVID-19 pandemic in Mexico

09 Jul 2022 - Biomed Central

While suicide rates in high- and middle-income countries appeared stable in the early stages of the pandemic, we know little about within-country variations. We sought to investigate the impact of COVID-19 on ...

Bullying and other risk factors related to adolescent suicidal behaviours in the Philippines: a look into the 2011 GSHS Survey

04 Jul 2022 - Biomed Central

The present study retrospectively examined gender differences in bullying and suicidal behaviour (ideation, plan, and attempts) as well as associations between selected risk factors and suicidal behaviour amon...

Suicide Attempt and Risk of Substance Use Disorders Among Female Youths

01 Jul 2022 - The JAMA Network

This cohort study assesses whether suicide attempts by females aged 8 to 19 years in Quebec, Canada, are associated with subsequent risk of substance use disorders later in life.

Prevalence and Correlates of Suicide and Nonsuicidal Self-injury in Children

01 Jul 2022 - The JAMA Network

This systematic review and meta-analysis assesses prevalence estimates of preadolescent self-injurious thoughts and behaviors, identifies correlates of these outcomes, and conducts head-to-head comparisons of preadolescent and adolescent self-injurious thoughts and behaviors in terms of associated characteristics.

[\[back to top\]](#)

Attention Deficit Hyperactivity Disorder

Editorial: The Epidemiology and Cognitive Characteristics of Attention-Deficit/Hyperactivity Disorder Depend on how Strictly the Disorder is Defined

05 Jul 2022 - Elsevier ScienceDirect

Publication date: Available online 5 July 2022Source: Journal of the American Academy of Child & Adolescent PsychiatryAuthor(s): Joseph Aloï, Leslie Hulvershorn

[\[back to top\]](#)

Obsessive Compulsive Disorder

Any overlap between orthorexia nervosa and obsessive–compulsive disorder in Lebanese adults? Results of a cross-sectional study and validation of the 12-item and 4-item obsessive–compulsive inventory (OCI-12 and OCI-4)

14 Jul 2022 - Biomed Central

Orthorexia Nervosa (ON), a compulsive obsession with vigorous eating, has increasingly caught researchers' attention. Although Orthorexia Nervosa has not been labeled an eating disorder, research about ON high...

[\[back to top\]](#)